

IJC Prelim 2 Paper 2 Suggested Solutions

Qn	Suggested Solutions
<p>1(i)</p>	$y = (\cos^{-1} x)^2$ $\frac{dy}{dx} = 2(\cos^{-1} x) \cdot \frac{-1}{\sqrt{1-x^2}}$ $\sqrt{1-x^2} \frac{dy}{dx} = -2\cos^{-1} x$ <p>diff wrt x,</p> $\sqrt{1-x^2} \frac{d^2y}{dx^2} + \frac{-2x}{2\sqrt{1-x^2}} \frac{dy}{dx} = -2 \left(\frac{-1}{\sqrt{1-x^2}} \right)$ $(1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} = 2 \quad (\text{shown})$ <p>Alternative</p> $y = (\cos^{-1} x)^2$ $\frac{dy}{dx} = 2(\cos^{-1} x) \cdot \frac{-1}{\sqrt{1-x^2}}$ <p>diff wrt x,</p> $\frac{d^2y}{dx^2} = 2 \cdot \frac{-1}{\sqrt{1-x^2}} \cdot \frac{-1}{\sqrt{1-x^2}} + (-2) \cdot (\cos^{-1} x) \cdot \frac{-1}{2} (1-x^2)^{-\frac{3}{2}} (-2x)$ $\frac{d^2y}{dx^2} = \frac{2}{(1-x^2)} + (-2x) \cdot (\cos^{-1} x) \cdot \frac{1}{(1-x^2)\sqrt{1-x^2}}$ $(1-x^2) \frac{d^2y}{dx^2} = 2 + (-2) \cdot (\cos^{-1} x) \cdot \frac{x}{\sqrt{1-x^2}}$ $(1-x^2) \frac{d^2y}{dx^2} = 2 + x \frac{dy}{dx}$ $(1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} = 2 \quad (\text{shown})$
<p>(ii)</p>	<p>diff wrt x,</p> $(1-x^2) \frac{d^3y}{dx^3} + (-2x) \frac{d^2y}{dx^2} - x \frac{d^2y}{dx^2} - \frac{dy}{dx} = 0$ $(1-x^2) \frac{d^3y}{dx^3} - (3x) \frac{d^2y}{dx^2} - \frac{dy}{dx} = 0$

	<p>when $x = 0$, $y = \left(\frac{\pi}{2}\right)^2$, $\frac{dy}{dx} = -\pi$, $\frac{d^2y}{dx^2} = 2$, $\frac{d^3y}{dx^3} = -\pi$,</p> $y = \left(\frac{\pi}{2}\right)^2 + (-\pi)x + \frac{2}{2!}x^2 + \frac{-\pi}{3!}x^3 + \dots$ $= \frac{\pi^2}{4} - \pi x + x^2 - \frac{\pi}{6}x^3 + \dots$
(iii)	$e^{(\cos^{-1}x)^2} = e^{\frac{\pi^2}{4} - \pi x + x^2 - \frac{\pi}{6}x^3} = e^{\frac{\pi^2}{4}} \cdot e^{-\pi x + x^2 - \frac{\pi}{6}x^3}$ $= e^{\frac{\pi^2}{4}} \cdot \left[1 + \left(-\pi x + x^2 - \frac{\pi}{6}x^3\right) + \frac{1}{2} \left(-\pi x + x^2 - \frac{\pi}{6}x^3\right)^2 + \dots \right]$ $= e^{\frac{\pi^2}{4}} \cdot \left[1 + \left(-\pi x + x^2 - \frac{\pi}{6}x^3\right) + \frac{1}{2}(-\pi x)^2 \dots \right]$ $= e^{\frac{\pi^2}{4}} \cdot \left[1 - \pi x + x^2 - \frac{\pi}{6}x^3 + \frac{1}{2}\pi^2 x^2 \dots \right]$ $= e^{\frac{\pi^2}{4}} \cdot \left[1 - \pi x + \left(1 + \frac{1}{2}\pi^2\right)x^2 + \dots \right]$ <p>$\therefore a = -\pi$ and $b = 1 + \frac{1}{2}\pi^2$</p>

Qn	Suggested Solutions
2	$y = vx$ $\frac{dy}{dx} = v + x \frac{dv}{dx}$ $x \frac{dy}{dx} - y = x(x - y)$ $x \left(v + x \frac{dv}{dx} \right) - vx = x(x - vx)$ $v + x \frac{dv}{dx} - v = x - vx$ $x \frac{dv}{dx} = x(1 - v)$ $\frac{dv}{dx} = 1 - v$

$$\Rightarrow \int \frac{1}{1-v} dv = \int dx$$


$$-\ln|1-v| = x + C$$

$$|1-v| = e^{-x-C}$$

$$1-v = \pm e^{-C} e^{-x}$$

$$v = 1 - Ae^{-x}$$

$$y = x(1 - Ae^{-x}) \quad (\text{shown})$$



At the stationary point, $\frac{dy}{dx} = 0$.

Hence, equation of locus is: $x(0) - y = x(x - y)$

$$y = -x(x - y)$$

$$y = -x^2 + xy$$

$$\text{i.e., } y = \frac{x^2}{x-1}$$

Qn	Suggested Solutions
3(i)	<p>Let $y = \frac{5}{(x-4)^2}$.</p> $(x-4)^2 = \frac{5}{y}$ $x-4 = \pm \sqrt{\frac{5}{y}}$ $x = \pm \sqrt{\frac{5}{y}} + 4 \quad (\text{reject positive sq root } \because x < 4)$ $x = 4 - \sqrt{\frac{5}{y}}$ $\therefore f^{-1}(x) = 4 - \sqrt{\frac{5}{x}}$ $D_{f^{-1}} = R_f = (0, \infty)$

(iii)	
(iv)	<p>Solve $f(x) = x$.</p> $\frac{5}{(x-4)^2} = x$ $5 = x(x-4)^2$ $5 = x(x^2 - 8x + 16)$ $x^3 - 8x^2 + 16x - 5 = 0 \text{ (shown)}$ $x^3 - 8x^2 + 16x - 5 = 0$ $(x-5)(x^2 - 3x + 1) = 0$ $x = 5, x = \frac{3 \pm \sqrt{9-4}}{2}$ $x = 5, x = \frac{3 + \sqrt{5}}{2} \text{ or } x = \frac{3 - \sqrt{5}}{2}$ <p>(rej as $x < 4$)</p>

4(i)	$\vec{OA} = -6\mathbf{i} - 3\mathbf{j} = \begin{pmatrix} -6 \\ -3 \\ 0 \end{pmatrix}, \vec{OC} = -\vec{OA} = \begin{pmatrix} 6 \\ 3 \\ 0 \end{pmatrix}, \vec{OV} = \begin{pmatrix} 0 \\ 0 \\ 6 \end{pmatrix}$ $\therefore \vec{OM} = \frac{1}{2} \left[\begin{pmatrix} 6 \\ 3 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ 6 \end{pmatrix} \right] = \begin{pmatrix} 3 \\ \frac{3}{2} \\ 3 \end{pmatrix}$ $\therefore \vec{AM} = \begin{pmatrix} 3 \\ \frac{3}{2} \\ 3 \end{pmatrix} - \begin{pmatrix} -6 \\ -3 \\ 0 \end{pmatrix} = \begin{pmatrix} 9 \\ \frac{9}{2} \\ 3 \end{pmatrix} = \frac{3}{2} \begin{pmatrix} 6 \\ 3 \\ 2 \end{pmatrix}$ <p>Hence, equation of the line AM is $\mathbf{r} = \begin{pmatrix} -6 \\ -3 \\ 0 \end{pmatrix} + t \begin{pmatrix} 6 \\ 3 \\ 2 \end{pmatrix}, t \in \mathbb{R}$</p>
------	---

(ii)

$$\vec{AB} = \begin{pmatrix} 12 \\ 0 \\ 0 \end{pmatrix}$$

Length of projection of \vec{AB} onto the line AM ,

$$AN = \frac{\left| \vec{AB} \cdot \begin{pmatrix} 6 \\ 3 \\ 2 \end{pmatrix} \right|}{\sqrt{6^2 + 3^2 + 2^2}}$$

$$= \frac{\left| \begin{pmatrix} 12 \\ 0 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} 6 \\ 3 \\ 2 \end{pmatrix} \right|}{7} = \frac{72}{7}$$

Perpendicular distance from P to the line AM

$$= \sqrt{(AB)^2 - (AN)^2}$$

$$= \sqrt{12^2 - \left(\frac{72}{7}\right)^2} = 6.18 \text{ (3 s.f.)}$$

Alternative Method

Perpendicular distance from P to the line AM

$$= \frac{\left| \vec{AB} \times \begin{pmatrix} 6 \\ 3 \\ 2 \end{pmatrix} \right|}{\sqrt{6^2 + 3^2 + 2^2}} = \frac{\left| \begin{pmatrix} 12 \\ 0 \\ 0 \end{pmatrix} \times \begin{pmatrix} 6 \\ 3 \\ 2 \end{pmatrix} \right|}{\sqrt{49}}$$

$$= \frac{\left| \begin{pmatrix} 0 \\ -2 \\ 3 \end{pmatrix} \right|}{7} = \frac{12\sqrt{4+9}}{7}$$

$$= 6.18 \text{ (3 s.f.)}$$

(iii)

A normal vector to the plane AMB

$$= \begin{pmatrix} 12 \\ 0 \\ 0 \end{pmatrix} \times \begin{pmatrix} 6 \\ 3 \\ 2 \end{pmatrix} = 12 \begin{pmatrix} 0 \\ -2 \\ 3 \end{pmatrix}$$

$$\vec{DV} = \begin{pmatrix} 0 \\ 0 \\ 6 \end{pmatrix} - \begin{pmatrix} -6 \\ 3 \\ 0 \end{pmatrix} = \begin{pmatrix} 6 \\ -3 \\ 6 \end{pmatrix} = 3 \begin{pmatrix} 2 \\ -1 \\ 2 \end{pmatrix}$$

Let θ be that angle betw. the line DV and plane AMB .

$$\sin \theta = \frac{\left| \begin{pmatrix} 2 \\ -1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 0 \\ -2 \\ 3 \end{pmatrix} \right|}{\sqrt{4+1+4}\sqrt{4+9}}$$

$$= \frac{2+6}{3\sqrt{13}}$$

$\theta = 47.7^\circ$ (1 d.p.)

(iv) If the 3 planes AMB , AMD and Π do not have a common point, the line AM is parallel to Π but does not lie in Π .

$$\therefore \begin{pmatrix} 6 \\ 3 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 4 \\ a \end{pmatrix} = 0$$

$$\Rightarrow -6 + 12 + 2a = 0$$

$$\Rightarrow a = -3$$

Note that $\begin{pmatrix} -6 \\ -3 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 4 \\ a \end{pmatrix} = 6 - 12 \neq 4,$

Therefore point A does not lie in Π .
Hence the line AM does not lie in Π .

Qn Suggested Solutions

5 Let X be the random variable denoting the volume of hot chocolate.

$$E(X) = 55, \quad \text{Var}(X) = 10^2$$

By Central Limit Theorem,

$$\bar{X} \sim N\left(55, \frac{10^2}{n}\right) \text{ approximately.}$$

$$P(\bar{X} > 54) > 0.77$$

$$P\left(Z > \frac{54 - 55}{10/\sqrt{n}}\right) > 0.77$$

$$P\left(Z < \frac{-\sqrt{n}}{10}\right) < 0.23$$

$$\frac{\sqrt{n}}{10} > 0.7388$$

$$\sqrt{n} > 10(0.7388) = 7.388$$

$$n > 54.59$$

\therefore the least $n = 55$

Qn	Suggested Solutions
6(i)	<ul style="list-style-type: none"> Obtain a list of households, in order of surnames in that particular constituency. Randomly select a starting point in the first 20 households on the list and thereafter select every 20th household on the list to be interviewed to get their responses.
6(iii)	<p>A better sampling method is <u>stratified sampling</u>.</p> <p>Obtain a list of all households and divide the households according to the different types of housing. Select a random sample from each stratum such that the sample size is proportional to the relative size of each type of housing.</p> <p>Thus, the sample obtained is more representative of the population.</p>


Qn	Suggested Solutions
7(i)	<p>Let X be the random variable denoting the number of visitors to the blog in two days.</p> <p>$X \sim \text{Po}(6.4)$</p> $P(5 \leq X < 10)$ $= P(X \leq 9) - P(X \leq 4)$ $= 0.650729 = 0.651 \text{ (3sf)}$
(ii)	<p>Let Y be the random variable denoting the number of visitors to the blog in a week.</p> <p>$Y \sim \text{Po}(22.4)$</p> <p>Since $\lambda = 22.4 > 10$</p> <p>$Y \sim N(22.4, 22.4)$ approx</p> $P(Y > 30) = P(Y > 30.5) \text{ with cc}$ $= 0.043500 = 0.0435 \text{ (3 sf)}$
(iii)	<p>The mean no. of visitors may not be the same for each day, e.g. weekends or during school holidays, mean no. of visitors may be higher.</p>

Qn	Suggested Solutions
8	
(i)	<p>no. of ways the married couple seated together = $2! \times 6! = 1440$</p> <p>Total no. of arrangement, without restriction = $7!$</p> $P(\text{no. of ways the married couple seated together}) = 1440/7! = 2/7$
(ii)	<p>$P(\text{all women sat together} \mid \text{couple sat together})$</p> $= \frac{P(\text{all women sat together and couple sat together})}{P(\text{couple sat together})}$ $= \frac{(6)3!2!}{6!2!}$ $= \frac{1}{20}$

(iii)	<p>no. of ways men and women alternate $= 4! \times 3! = 144$</p> <p>$P(\text{men and women alternate}) = \frac{144}{7!} = \frac{1}{35}$</p>
(iv)	<p>no. of ways to arrange around a table with no restriction $= (7-1)! = 720$</p> <p>no. of ways to arrange around a table with one particular woman must sit between two men $= {}^3P_2 \times (5-1)! = 144$</p> <p>$P(\text{no. of ways to arrange around a table with one particular woman must sit between two men}) = 144/720 = 1/5$</p>

Qn	Suggested Solutions
9(i)	<p>Let X be the random variable denoting the number of students who are awarded distinction out of 23 students. $X \sim B(23, 0.2)$ $E(X) = np = 4.6$</p>
(ii)	<p>Let Y be the random variable denoting the number of students who are awarded distinction out of n students. $Y \sim B(n, 0.2)$</p> <p>$P(Y \geq 9) > 0.7$ $1 - P(Y \leq 8) > 0.7$ $P(Y \leq 8) < 0.3$ From GC, when $n = 50$, $P(Y \leq 8) = 0.30733$ $n = 51$, $P(Y \leq 8) = 0.28395$</p> <p>\therefore smallest $n = 51$</p>
(iii)	<p>Let S be the random variable denoting the number of students who pass out of 60 students. $S \sim B(60, 0.93)$ $n = 60$ is large, $np = 55.8 > 5$, $nq = 4.2 < 5$</p> <p>$S' \sim B(60, 0.07)$ Since $n = 60$ is large, $np = 4.2 < 5$, $S' \sim \text{Po}(4.2)$ approx.</p> <p>$P(S > 55) = P(S' < 5)$ $= P(S' \leq 4) = 0.58982 = 0.590$</p>

Qn	Suggested Solutions
10	<p>Let A and C be the random variables denoting the mass of a randomly chosen almond cookie and chocolate chip cookie respectively.</p> $A \sim N(32, 3.0^2)$ $C \sim N(28, 1.8^2)$ <p>(i) $E(C_1 + C_2 + C_3 + C_4) = (4)(28) = 112$ $\text{Var}(C_1 + C_2 + C_3 + C_4) = (4)(1.8^2)$</p> $C_1 + C_2 + C_3 + C_4 \sim N(112, 12.96)$ $P(100 < C_1 + C_2 + C_3 + C_4 < 120)$ $= 0.9864368$ $= 0.986(3sf)$
(ii)	$E(C_1 + C_2 + C_3 + C_4 - 2A) = 4E(C) - 2E(A) = (4)(28) - 2(32) = 48$ $\text{Var}(C_1 + C_2 + C_3 + C_4 - 2A) = 4\text{Var}(C) + 2^2\text{Var}(A) = (4)(1.8^2) + 2^2(3.0^2) = 48.96$ $C_1 + C_2 + C_3 + C_4 - 2A \sim N(48, 48.96)$ $P(-50 < C_1 + C_2 + C_3 + C_4 - 2A < 50)$ $= 0.6124961464 = 0.612(3sf)$
(iii)	<p>Let $X = \frac{6}{100}(A_1 + A_2 + A_3 + A_4) + \frac{5}{100}(C_1 + C_2 + C_3 + C_4)$ $- \frac{5}{100}(C_{11} + C_{12} + \dots + C_{20})$</p> $E(X) = \frac{6}{100}(4)E(A) - \frac{5}{100}(6)E(C) = -0.72$ $\text{Var}(X) = \left(\frac{6}{100}\right)^2 (4)\text{Var}(A) + \left(\frac{5}{100}\right)^2 (14)\text{Var}(C) = 0.243$ $X \sim N(-0.72, 0.243)$ $P(X > 0) = 0.0720635555 = 0.0721(3sf)$

Qn	Suggested Solutions
11	
(i)	 <p data-bbox="259 735 1396 840">From the scatter plot, over time, there is an increase in the rate of increase of the no. of mosquitoes. Hence it shows a curvilinear relationship between t and x. A linear model will not be appropriate.</p>
(ii)	<p data-bbox="235 913 341 945">$x = ae^{bt}$</p> <p data-bbox="235 955 414 987">$\ln x = \ln a + bt$</p> <p data-bbox="235 1050 365 1081">From GC,</p> <p data-bbox="235 1092 479 1123">$\ln a = 3.114769038$</p> <p data-bbox="235 1134 625 1165">$b = 0.3555454706 = 0.356$ (3sf)</p> <p data-bbox="235 1228 479 1260">$\ln x = 3.11 + 0.356 t$</p> <p data-bbox="235 1312 755 1354">$\therefore a = e^{3.114769038} = 22.52822659 = 22.5$ (3sf)</p>
(iii)	<p data-bbox="235 1459 397 1491">When $t = 30$,</p> <p data-bbox="592 1501 828 1533">$\ln x = 13.78113316$</p> <p data-bbox="592 1543 738 1575">$x = 966207$</p>
(iv)	<p data-bbox="235 1669 1201 1711">The calculated value for x is unreliable since $t = 30$ is outside the data range of t.</p>

Qn	Suggested Solutions
12(i)	<p>unbiased estimate of population mean,</p> $\bar{x} = \frac{\sum(x-1200)}{120} + 1200 = \frac{-60}{120} + 1200 = 1199.5$ <p>unbiased estimate of population variance,</p> $s^2 = \frac{1}{119} \left[2014 - \frac{(-60)^2}{120} \right] \approx 16.672 = 16.7 \text{ (3 s.f.)}$
(ii)	<p>Let μ hrs be the population mean lifetime of the bulbs.</p> <p>$H_0 : \mu = 1200$</p> <p>$H_1 : \mu < 1200$ (manufacturer's claim invalid)</p> <p>Level of significance: 10%</p> <p>Since population variance is unknown and sample size $n = 120$ is large, by Central Limit Theorem, $\bar{X} \sim N\left(\mu, \frac{s^2}{n}\right)$ approximately.</p> <p>Under H_0, test statistics, $Z = \frac{\bar{X} - 1200}{s/\sqrt{120}} \sim N(0,1)$ approximately</p> <p>From GC, $p\text{-value} = 0.089878 = 0.0899$ (3 s.f.)</p> <p>Since $p\text{-value} = 0.0899 (< 0.1)$, reject H_0 and conclude that there is sufficient evidence at the 10% level of significance that the manufacturer's claim is not valid.</p>
(iii)	<p>Assume that the lifetime of the light bulbs produced by the improved process follows a normal distribution.</p> <p>$H_0 : \mu = 1200$</p> <p>$H_1 : \mu < 1200$</p> <p>Level of significance = 5%</p> <p>Under H_0, test statistic: $T = \frac{\bar{X} - 1200}{S/\sqrt{20}} \sim t_{(19)}$</p> <p>Critical region : Reject H_0 if $t \leq -1.72913$.</p> <p>Given sample mean = k and sample std. dev. = 9.8</p> <p>\therefore unbiased estimate of population variance,</p> $s^2 = \frac{n}{n-1} \times (\text{sample variance}) = \frac{20}{19} (9.8)^2 = 101.095$ <p>For the test to indicate that the manufacturer's claim is valid for this improved process, we</p>

do not reject H_0 .

$$\text{Hence, } \frac{k-1200}{\frac{\sqrt{101.095}}{\sqrt{20}}} > -1.72913$$

$$k > 1196.11243$$

The least possible value of k is 1196.113 (or 1196.112)
(3 d.p.)